

International Journal of Population Data Science

Journal Website: www.ijpds.org



The Kids' Environment and Health Cohort: a novel administrative data resource for research on the environmental determinants of child health in England

Selin Akaraci¹, Alison Macfarlane², Amal Rammah¹, Emily Courtin³, Faith Miller¹, Jessica Mitchell¹, Joana Cruz¹, Matthew Lilliman¹, Niloofar Shoari¹, Samantha Hajna⁴, Steven Cummins⁵, Vahe Nafilyan⁶, and Pia Hardelid¹

¹University College London

²City University of London

³The London School of Economics and Political Science

⁴Brock University

⁵London School of Hygiene and Tropical Medicine

⁶Office for National Statistics

Objectives

Through the Multidisciplinary Ecosystem to study Lifecourse Determinants and Prevention of Early-onset Burdensome Multimorbidity (MELD-B) project, we established the SAIL MELD-B e-cohort (SMC) and the SAIL MELD-B Young adults e-cohort (SMYC), with the aim to enhance the understanding of 'burdensomeness' in individuals living with multimorbidity, including identifying new clusters of burdensome indicators, exploring early life risk factors and modelling potential preventative scenarios.

Approach

We use routinely-collected anonymised linked demographic, health and administrative data sources available within the SAIL Databank to define SMC and SMYC. These cohorts were developed using a reproducible, maintainable, methodological pipeline that allows for dynamic updates as data coverage expands. The pipeline efficiently processes new burdensomeness concepts, facilitating the extraction of relevant records associated with the concepts identified for use in the SMC and SMYC.

Results

SMC and SMYC comprises of 5,180,602 and 896,155 individuals registered with a Welsh General Practice at any time between 1st January 2000 and 31st December 2022 respectively. Analysis of primary and secondary care health data reveals that the most common conditions in SMC were depression (21.6%), anxiety (21.1%), asthma (17.5%), hypertension (16.2%), and atopic eczema (14.1%). In SMYC, the most common conditions were atopic eczema (21.2%), asthma (11.6%), anxiety (6.0%), deafness (4.6%), and depression (4.3%).

Conclusions and Implications

SMC and SMYC provide two generalisable population samples, which can be used to address various research questions across MELD-B. The adaptability of the methodological pipeline allows cohort curation to be repurposed for other projects accessing population-scale data sources and trusted research environments.

